




ELSEVIER

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/ecolmodel

How important for wintering is the location of breeding areas in fragmented landscapes? Maximum entropy niche-based modelling of seasonal changes in little bustard (*Tetrax tetrax*) distribution

Susana Suárez-Seoane^{a,*}, Eladio L. García de la Morena^b, Manuel B. Morales Prieto^b, Patrick E. Osborne^c, Eduardo de Juana^d

^a Area de Ecología, Facultad de Ciencias Biológicas y Ambientales, Universidad de León, Campus de Vegazana s/n, 24071 León, Spain

^b Departamento Interuniversitario de Ecología, Facultad de Ciencias (Biología), Universidad Autónoma de Madrid, 28049 Madrid, Spain

^c Centre for Environmental Sciences, School of Civil Engineering and the Environment, University of Southampton, Highfield, Southampton SO17 1BJ, UK

^d Departamento de Biología Animal I, Facultad de Biología, Universidad Complutense de Madrid, 28040 Madrid, Spain

ARTICLE INFO

Article history:

Received 27 August 2007

Received in revised form

16 July 2008

Accepted 17 July 2008

Keywords:

Connectedness

Habitat and climate preferences

MaxEnt

METEOSAT

Niche width

ABSTRACT

The effects of habitat fragmentation on species may change seasonally mainly due to variations in resource availability and biotic interactions. In critical periods, such as winter, when the importance of intraspecific competition diminish, species may relax their environmental requirements widening their ecological niche to exploit trophic resources more efficiently in comparison with spring. Those variations in niche width may implicate seasonal expansions/retractions in species distribution. In this sense, an integrated knowledge on the spatial arrangement of breeding and wintering suitable patches is essential to infer seasonal movements (migratory connectivity). This paper shows that little bustard environmental preferences were more predictable and complex (controlled by a larger number of environmental factors) in spring than in winter, when potential distribution and ecological niche width were slightly larger. In spring, habitat variables (i.e. dry crops, pasturelands and altitude) ruled species' distribution; while, winter pattern was driven by mixed criteria, based on both habitat and climate (i.e. dry crops, wastelands and winter rainfall). Suitable patches were more connected across spatial scales in winter than in spring, i.e. landscape was perceived as less fragmented. The overlap between potential breeding and wintering distribution areas was high. In fact, most of the predicted wintering areas coincided or showed high connectedness with predicted breeding patches. Conversely, there were significant breeding patches that were predicted as basically unsuitable, showing little connectedness with potential winter areas. Spring habitat is a better predictor of little bustard's wintering range than vice versa, which has clear management implications (preserving breeding sites closer to wintering areas ensures the conservation of a larger proportion of the total distribution range). This is an example of how predictive large-scale modeling procedures can contribute to the optimization of land management aimed at species conservation.

© 2008 Published by Elsevier B.V.

* Corresponding author. Tel.: +34 987 291565.

E-mail address: s.seoane@unileon.es (S. Suárez-Seoane).

0304-3800/\$ – see front matter © 2008 Published by Elsevier B.V.

doi:10.1016/j.ecolmodel.2008.07.035

1. Introduction

The fragmentation of a habitat into discontinuous patches negatively affects population recruitment (Robinson et al., 1995; Smith and Hellmann, 2002), survival (Harris, 1984) and movement (Shirley, 2006) of terrestrial animal species. In poorly connected landscapes, where individuals have to move across the matrix to reach adequate pieces of habitat for different purposes (i.e. foraging, reproduction, dispersion and predator avoidance), the fitness cost of movement (Brooker et al., 1999) becomes higher than in continuous landscapes. This fact consequently influences the dynamics, spatial structure and persistence of populations (Turchin, 1991). Nevertheless, the importance of fragmentation for species may change through time, mainly seasonally, individual movements reflecting variations in resource availability (Blake and Loiselle, 1991) and biotic interactions. In this context, linking breeding and non-breeding populations (i.e. migratory connectivity) to infer movement patterns between seasons is one of the ultimate goals of animal movement studies in ecology (Rubenstein and Hobson, 2004).

As a result of the European Common Agricultural Policy, traditional agri-systems in southern Europe are particularly vulnerable to fragmentation due to management intensification in productive areas and land abandonment in marginal ones (Pain and Pienkowski, 1997). These systems are known to host a considerable diversity of birds and other taxa, most of which are currently experiencing marked population declines (Robinson and Sutherland, 2002; Sanderson et al., 2005). Among the valuable avian species (Suárez et al., 1997) held by these systems, the little bustard *Tetrax tetrax* is one of most seriously threatened by land use changes (Wolff, 2001; Wolff et al., 2002; García et al., 2007), disappearing from many European countries during recent decades (Schulz, 1985; Goriup, 1994; Del Hoyo et al., 1996). This Palaearctic, medium-sized steppe bird, from the *Otididae* family, is currently classified as ‘Near Threatened’ (Collar et al., 1994) in the world and ‘Vulnerable’ in Europe (BirdLife International, 2004), including Spain (García de la Morena et al., 2004b). Although Iberian little bustard populations have been regarded as sedentary or dispersive (Cramp and Simmons, 1980), many of them can actually be considered as migratory or partially migratory since they completely, or partly, disappear from their breeding grounds, performing regular medium or long distance movements and congregating in certain wintering sites (García de la Morena et al., 2004a, 2006). In Madrid region (the study area), it is considered a resident species (Díaz et al., 1994; García de la Morena, 2002), although it exhibits a partial migratory behaviour, as suggested by recent radiotracking data (with some birds leaving the region during the non-breeding season; own unpubl. data). During the spring (breeding season), birds do not move much, spending most of time in their territories (Schulz, 1985; Jiguet, 2001) but, from late summer through the autumn and winter, they gather in flocks and disperse from spring areas to exploit food resources in different zones, a common behaviour in most Iberian steppe birds (Suárez et al., 1997). As found in some of these species (Morales et al., 2000; Alonso et al., 2001), little bustards display interannual fidelity to certain wintering sites, where they stay for a variable period of time before returning to their breed-

ing territories (García de la Morena et al., 2004b; own unpubl. data).

Habitat preferences and distribution of little bustards in spring have been extensively explored at both local (Martínez, 1994, 1998; Salamolard and Moreau, 1999; Wolff et al., 2001; Morales et al., 2005, in press; Traba et al., 2008) and regional scale (Suárez-Seoane et al., 2002, 2004; Osborne and Suárez-Seoane, 2002; García et al., 2007). However, few authors have focused on winter season (Leitão and Costa, 2001; Silva et al., 2004; García de la Morena et al., 2006, 2007), even if this knowledge is essential for understanding the species’ biological cycle, as well as in the design of adequate conservation strategies (Rappole and McDonald, 1994; Sherry and Holmes, 1996). In the Mediterranean region, as in other temperate and seasonally regulated areas, winter is a critical period for birds (Tellería et al., 1988), both at individual and population levels (Wiens, 1989; Newton, 1998), since availability in food and other resources decrease and may vary considerably in space and time. As a result, the distribution and abundance of wintering little bustards is closely dependent on the local variation of those resources (mainly provided by extensive cereal farmlands), which they must track actively (Wolff, 2001). During this limiting period, birds may therefore respond by relaxing the requirements associated to certain dimensions of their ecological niche to exploit more efficiently a larger amount of trophic resources, thus allowing the species’ survival (Hutchinson, 1957) in a wider potential distribution range. As a consequence of this niche expansion, wintering populations are expected to become more heterogeneous in their environmental preferences, according to the niche variation hypothesis (Van Valen, 1965; Bolnick et al., 2007), which predicts that each individual might continue to use a narrow range of resources but diverge from its conspecific competitors to minimize resource use overlap and competition. At the same time, intraspecific interactions would become less intense than interspecific relationships (Morin and Chuine, 2006), such as competition or predation (birds have to aggregate in flocks as a defense strategy against predators), in comparison with spring, when the relevance of territorialism and sexual behavior is higher. As a consequence of this shift in the importance of inter/intraspecific interactions, the relevance of certain niche dimensions (e.g. climate conditions) would be relatively higher than in spring.

To approach seasonal variations in species’ niche dimensions and their influence in spatial distribution, we took advantage of using the Maximum Entropy Modelling (MaxEnt; Phillips, 2005; Phillips et al., 2006). This novel technique provides a general-purpose machine learning method whose performance has been evaluated as one of the best when compared to other modelling distribution methods, particularly at small sample sizes (Elith et al., 2006; Hernández et al., 2006; Pearce and Boyce, 2006; Pearson et al., 2007). The following are among the main reasons to be used in this study: (1) it is an envelope-method specifically applied on presence-only data (the link between absences and habitat suitability may be confusing); (2) it has a good ability to fit complex functions between response and predictor variables; and (3) model selection and fitted models are not too complex, being similar in expressiveness to a GLM or GAM. As other niche-based models, MaxEnt describes suitability in

ecological space, which is projected into geographical space. Therefore, areas that satisfy the conditions of the species' fundamental niche represent its potential distribution, whereas the geographical areas that it actually inhabits constitute its realized distribution (Phillips et al., 2006). Nevertheless, Araújo and Guisan (2006) highlight several difficulties of using Hutchinson's (1957) concepts of fundamental and realized niches in species distribution modelling at large scale. In this sense, it is key to clarify the difference between "niche" and "area of distribution" (Soberon, 2007), distinguishing between niches as: (1) habitat or function; (2) characterized at local (when behavior and physiology are important) or at larger spatial extents (when distributional limits matter); and (3) defined by interactive variables such as resources (i.e. "Eltonian niches"), which can mainly be measured at local scales, or defined by non-interactive variables such as abiotic conditions (i.e. "Grinnellian niches"), relevant to understand coarse-scale ecological and geographical properties of species. Both classes of niches are relevant to understand species' distribution, but the Grinnellian ones are more appropriate at lower spatial resolutions and wider extensions, at which distributions are typically defined, as it is the case of the present study. Concerning the relevant niche dimensions to be explored at those large scales, climatic preferences can be used to predict areas where species could occur, since climate is the major driving factor of species' distribution (Thuiller et al., 2005), although Broennimann et al. (2007) showed that some climatic factors may only be indirectly related to niche shifts. Therefore, other non-climatic factors such as vegetation, geology or disturbances may become highly relevant to be included in the models (e.g. Acevedo et al., 2007), as they determine species presence at finer scales.

According to this background, the main objective of this paper is to analyse how little bustard's perceive landscape fragmentation in two critical periods (spring and winter), focusing on seasonal spatial variations in environmental (habitat and climate) preferences, niche width and connectedness of the predicted distribution patterns. In particular, we are concerned with the assessment of the importance of the spatial arrangement of suitable breeding patches for the selection of wintering sites and vice versa, which may influence seasonal movements across the landscape (and therefore migratory connectivity). More specifically, we explore the following hypotheses: (1) In winter, territorial behaviour becomes less important and resource availability decreases, therefore environmental preferences will relax and ecological niche will expand. This will allow birds to exploit the scarce resources across a broader range of conditions, reducing intraspecific competition. (2) Analogously, landscape suitability and its connectedness will be lower in spring, when birds will show more restricted environmental preferences as they will look for more particular resources/conditions to establish their breeding territories under higher conspecific competition. (3) Finally, the roles played by both habitat and climate factors on the species' distribution will change across seasons; in particular, we predict a greater importance of climate variables in determining wintering patterns. The results of this paper have important potential implications for improving the design of conservation measures addressed to protect this species by identifying key areas which ensure its connectivity across the

year at a regional scale, the most pertinent for environmental and agricultural policies (Rounsevell et al., 2003).

2. Materials and methods

2.1. Study area

Madrid region is located in Central Spain (41°10'N, 39°53'N, 4°35'W, 3°03'W) and occupies about 8000 km². It can be broadly divided into two geographical units: (1) the mountains of the Spanish Central Range, which run along the north-east to south-west limit of the region, over 900–1,000 m.a.s.l. and (2) the plains which extend over the rest of the region as part of the Spanish Southern Plateau, between 430 and 900 m.a.s.l. (Fig. 1). This unit is, in fact, formed by a mosaic of different habitats, including extensive pastures, shrubs, forests and urban areas, although extensive cereal cultivation (with interspersed olive groves and vineyards) is dominant. It also holds the highest human population densities within the region (about 5.8 million people are concentrated in Madrid city and its surroundings). In spring, little bustards are widely distributed over this second unit (Díaz et al., 1994; García de la Morena et al., 2001), their presence and abundance being positively correlated with landscape heterogeneity and proportion of arable surface (Morales et al., 2005). In winter, the species principally occupies extensive cereal farms at the northeast and south of the region, but also uses other habitats, such as irrigated alfalfa (*Medicago sativa*) fields, open broom formations dominated by *Retama sphaerocarpa* or other shrublands (García de la Morena, 2002; García de la Morena et al., 2007).

2.2. Bird data

Wintering data were obtained in early February (13th–16th) 2003 and late January (23th–26th) 2004 during surveys carried out by car over the species' regional winter range. We used the look-see counting and mapping method (Bibby et al., 2000) adapted for a large area (see García de la Morena et al., 2007 for survey details). Spring records corresponded only to breeding adult males and were gathered from an unpublished report on species distribution and population estimates in Madrid region (García de la Morena et al., 2001). Counts were carried out from the 18th April to the 1st June of 2000, coinciding with the peak of male sexual display (Schulz, 1985; Martínez, 1994). This survey followed the standardised census methodology recommended for breeding males by most species' experts (García de la Morena et al., 2006). For later analyses, all locations were overlapped in a grid at 1 km² (pixel size) to match the resolution of the predictors used for model development.

2.3. Environmental predictors

Habitat variables (topography, human disturbances and land covers) were considered as "static" or "non-variant" across the seasons during the study period (Table 1). For topographic features, a digital terrain model with a resolution of 200 m was built and then transformed into derived variables (mean altitude and topographic variability) calculated within a moving window of 1 km² (see Suárez-Seoane et al., 2002 for more




Fig. 1 – Location of the study area.

253 details). Human disturbance variables (density of towns and
 254 roads) were obtained from vector maps at 1:200,000 (Span-
 255 ish National Geographic Institute) and rasterised into grids
 256 of 200 m resolution, subsequently converted into quantita-
 257 tive variables at 1 km². Land cover classification was based

on the official 1:50,000 habitat cartography of the Madrid
 Region (CAM, 1998), updated in the field in 2004. Relevant
 variables (percentage of irrigated lands, dry crops, pasture-
 lands, olive trees and wastelands) were measured using the
 1 km² moving window and then log-transformed to reduce

258
 259
 260
 261
 262

Table 1 – List of habitat (topography, human disturbances and land covers) and climate predictors used for modelling seasonal species distribution in Madrid region

Code	Variable
Habitat variables	
Topography	
MDT	Mean altitude within a 5 × 5 array of 200 m pixels
TOPO10	Variation in altitude in a 5 × 5 pixel array of 200 m pixels, where altitude is measured to 10 m vertical resolution. Calculated as $TV_x = \frac{(n-1)/(p-1)}{\lambda}$ where n = no. of different altitude classes in the array, p = no. of pixels in the array (i.e. 25), and λ = vertical resolution
Human disturbances	
TOWNDENS	Proportion of 200 m pixels in a 5 × 5 array containing buildings or large built structures such as airfields
ROADDENS	Proportion of 200 m pixels in a 5 × 5 array containing roads
Land covers	
IRRIGPERC	Proportion of 200 m pixels in a 5 × 5 (1 km ²) array containing irrigated lands
DRYPERC	Proportion of 200 m pixels in a 5 × 5 (1 km ²) array containing dry croplands
PASTPERC	Proportion of 200 m pixels in a 5 × 5 (1 km ²) array containing pasturelands
OLIPERC	Proportion of 200 m pixels in a 5 × 5 (1 km ²) array containing mosaic of cereal and olive trees
WASTEPC	Proportion of 200 m pixels in a 5 × 5 (1 km ²) array containing wastelands (old fallows, abandoned crops and suburban areas)
Climate variables	
EVPs/EVPw	Mean potential evapotranspiration value (calculated from the values per each 10-day periods) for the whole of the season (spring/winter)
RADs/RADw	Mean net radiation value (radiation balance calculated from observed albedo, temperatures and cloudiness according the time of the year and the position for each 10-day period) for the whole of the season (spring/winter)
RAINWs/RAINWw	Mean rainfall value during the last winter for spring bird data, and mean value of this parameter during the current season for winter bird data

collinearity caused by the unit sum constraint (Aebischer et al., 1993). Operations were performed using IDRISI Kilimanjaro14.01 (Eastman, 2003).

Climatic variables were considered as “non-static” or “variant” between spring and winter (Table 1). For each season, we created three time-dependent predictors related to rainfall, net radiation and potential evapotranspiration (as an indicator of soil moisture) from the European Energy and Water Balance Monitoring System EWBMS products (Rosema, 1993; Rosema et al., 2001), which are derived from the METEOSAT satellite, with a frequency of 10-days and a resolution of 5 km. This dataset has been evaluated before for modelling the distribution of little bustards in Spain by Suárez-Seoane et al. (2004), showing very good results. In particular, we extracted: (1) mean values of potential evapotranspiration for each season of bird data collection (spring: mid-April to beginning of June 2000; winter: beginning of October to mid-February 2002–2003 and 2003–2004); (2) idem for net radiation; (3) mean rainfall during the winter season for winter bird data; and, mean rainfall during the latest winter for spring bird data, according to Morales et al. (2002), who have shown a positive link between great bustard *Otis tarda* breeding productivity and winter precipitation.

2.4. Statistical analyses

Firstly, we explored the dataset by means of a univariate analysis (Mann–Whitney U-test), which provides a general picture of the importance of each environmental variable for characterizing both the distribution and the ecological niche of little bustard in Madrid region across seasons.

Then, we ran a collection of models using the MaxEnt 2.2 method by Phillips (2005) and Phillips et al. (2006). The approach of MaxEnt is to find the probability distribution of maximum entropy (closest to the uniform) subject to the constraints imposed by the information available regarding the observed distribution of the species and the environmental conditions across the study area. The method assigns a probability of occurrence to each cell grid in this area. Because the sum of the probabilities must equal 1, each probability is typically extremely small, making model output difficult to interpret. We therefore present the MaxEnt output (model predictions) as cumulative probabilities, where the value of a given pixel is the sum of that pixel and all others with equal or lower probability, multiplied by 100 to give a percentage. A high value (close to 100%) at a particular pixel indicates that it is predicted to have suitable conditions for the species. Note that using presence-only data, it is generally not possible to calculate probabilities of presence; instead, outputs are relative likelihood of presence (Pearce and Boyce, 2006). Regularization values, which reduce over-fitting, were selected automatically by the program. The recommended default values were used for both the convergence threshold (10^{-5}) and maximum number of iterations (500). The selection of environmental variables was also carried out automatically, following default rules dependent on the number of presence records. The default is to include all variables, i.e. no selection.

Spatial models were built separately for each season (spring or winter) and group of predictors (Table 1): (1) habitat variables, (2) climate, and (3) all together. In each case, occurrence

locations (the dependent variable) were randomly partitioned into two sub-samples: 80% used as training data set and the remaining 20% reserved for testing the resulting models (partitioned models). In addition, for visual interpretation, the algorithm was run on the 100% of occurrence points (full models), taking advantage of all available data to provide the best estimates of the species’ potential distribution. The performance of both full and partitioned models was evaluated by means of an adaptation to presence-only data sets of the area under the receiver operating characteristic (ROC) curve (AUC; Beck and Shultz, 1986; Zweig and Campbell, 1993; Fielding and Bell, 1997). This measure can be interpreted as the probability that a presence site will be ranked above a random background site (by default MaxEnt generate a sample of 10,000 background “pseudo-absences” uniformly at random to represent the environmental conditions in the region) (Phillips et al., 2006; Phillips, 2008). A random ranking has a value of around 0.5, while a perfect ranking achieves the maximum possible AUC of 1.0. Models with an AUC value above 0.75 are considered as potentially useful (Elith et al., 2006).

As we are dealing with two occurrence datasets (winter and spring) of different size, we did a preliminary analysis (i.e. randomly selecting the spring sample size to be the same as that in the winter) to test whether this fact affected MaxEnt final models.

Continuous model outputs (cumulative probability values ranging from 0 to 100%) can be transformed into Boolean maps of suitable–unsuitable areas through the application of different thresholds or “cut-offs” (i.e. all pixels showing values above a selected threshold are reclassified as “1” and the remaining pixels as “0”). However, as highlighted by Phillips et al. (2006), determining the optimal threshold still remains a little explored topic when MaxEnt is applied. In fact, no general purpose rule has yet been developed but, in general terms, it must be considered that as larger thresholds are selected, commission errors will decrease, but omission errors will increase (Fielding and Bell, 1997; Hernández et al., 2006). In order to accomplish further comparative analyses, we decided to use three alternative thresholds with different degree of restriction. In particular, we selected the corresponding cumulative value for an omission error of 10% (which maintain a high proportion of presences correctly predicted when compared with “the lowest presence threshold”; see Pearson et al., 2007), 5% and 15% (note that omission error values range from 0 to 100%). Results were assessed through changes in connectedness, which was measured on the suitable patches identified on the Boolean maps by applying the different thresholds. We used three indices: (i) a patch cohesion index (COHESION), which is computed from the information contained in patch area and perimeter and increases as the patch becomes more aggregated; and, two types of contagion indices which show the frequency of adjacencies between the same patch type on the map (they will take the minimum value when the class is maximally disaggregated and the maximum when the class is maximally clumped): (ii) CLUMPY, which it is corrected from random, and (iii) PLADJ, which does not account for random effects. Analyses were executed in Fragstat 3.3 (McGarigal and Marks, 1995).

Finally, to look at the relationships between winter and spring patterns, we joint models in two different ways: (1)

We explored the advantages of including the final integrated spring model (continuous values) as a predictor to explain wintering distribution and vice versa. (2) Once the use of a particular threshold was decided, we combined Boolean integrated models for both seasons into a new layer showing wintering, breeding and resident areas.

3. Results

In winter, a total of 78 flock locations were gathered but, when these records were overlapped on the 1-km²-resolution grid, only 53 pixels could be used as presence records for modelling. In spring, 417 records of males were collected and transformed into 211 occupied 1-km² pixels. The preliminary evaluation of the effect of the different sampling size in modelling seasonal distribution showed no significant differences (in both AUC and spatial pattern) when we randomly equalled the pool of presences for spring and winter.

3.1. Seasonal changes in environmental preferences

Exploratory univariate analyses of the input data showed seasonal differences between spring and winter (Table 2). Winter locations were significantly characterised by lower seasonal values (mean and variation) of both net radiation and evapotranspiration, indicating that, in general terms, birds tolerated a narrower range of climatic conditions than in spring. However, in this season, birds selected areas where winter rainfall remained more constant around medium values, at least for the study period. The analysis also showed that wintering, as well as spring habitats, corresponds to extensive landscapes (i.e. little bustards preferred dry croplands, as compared to, for example, irrigated lands in both seasons). However, in winter, the birds used a higher proportion of wastelands and were found at a lower altitude than in spring, when they selected pasturelands located at a higher altitude, in the northern part of the province.

Table 3 – AUC-values for habitat, climate and integrated seasonal models (random AUC is 0.5)

	Spring	Winter
Habitat		
Training (test) sets	0.933 (0.914)	0.923 (0.871)
Full model	0.932	0.921
Climate		
Training (test) sets	0.837 (0.805)	0.848 (0.837)
Full model	0.838	0.863
Integrated models		
Training (test) sets	0.944 (0.920)	0.923 (0.871)
Full model	0.945	0.940

All AUC values are significant at $p < 0.001$.

In general, both spring and winter models showed a good performance and predictive capability, although some differences can be noticed (Table 3). Dealing with habitat and climate independently, the spring models had the highest performance (see AUC values of training and full models) and predictive capability (see AUC values of test sets) when explaining habitat, but showed poorer explanatory power for climate than winter models. When climate and habitat were considered together in the integrated models, the spring model still was the most explanatory and predictable for the training/test sets, but seasonal differences became not much important when full models were analysed (from 0.945 to 0.940). Table 4 shows that spring models were, in general, more complex because they explained the birds' distribution using a larger number of relevant variables (which greatly affected the models when they were dropped). In spring, the most significant variable for modelling the species' habitat preferences was the percentage of dry crops, followed by altitude and the percentage pasturelands. For the climate model, the three climate variables included were all highly relevant for model construction, particularly net radiation. When both sets of predictors were analysed together in the integrated models,

Table 2 – Comparison (means and standard deviations) between variables quantifying both spring and winter habitat for the little bustard in the study area

	Spring (n = 211)		Winter (n = 53)		Significance U-test
	X	S.D.	X	S.D.	
MDT	691.78	56.83	639.46	61.83	0.000
TOPO10	0.07	0.08	0.04	0.06	0.074
TOWNDENS	0.01	0.05	0.08	0.04	0.719
ROADDENS	0.07	0.11	0.07	0.11	0.786
IRRIGPERC	0.01	0.06	0.01	0.07	0.402
DRYPERC	0.54	0.24	0.80	0.27	0.632
PASTPERC	0.05	0.18	0.00	0.00	0.024
OLIPERC	0.02	0.11	0.01	0.04	0.693
WASTEPERC	0.01	0.05	0.07	0.19	0.002
EVPs/EVPw	28.49	1.18	5.84	0.27	0.000
RADs/RADw	51.02	1.86	10.18	0.41	0.000
RAINWs/RAINWw	8.90	0.80	17.44	1.14	0.000

The table shows the significance of the Mann-Whitney U-test between two independent samples (only significant values are in bold). Note that the climate data layers differed between seasons while those for the other variables did not.

Table 4 – Contribution of each environmental variable to the gain of the models

	Spring			Winter		
	Habitat	Climate	All	Habitat	Climate	All
MDT	18.261		10.000	3.846		0.858
TOPO10	2.174		1.739	2.137		0.429
TOWNDENS	0.870		0.435	0.427		0.429
ROADDENS	2.609		1.739	1.709		1.288
IRRIGPERC	2.174		1.304	0.855		0.858
DRYPERC	33.043		15.217	31.624		18.455
PASTPERC	10.870		7.826	0.427		0.429
OLIPERC	5.652		3.913	0.000		0.000
WASTEPCRC	2.174		1.739	7.692		5.579
EVP		19.535	0.870		0.730	0.000
RAD		32.558	1.304		31.387	2.146
RAINW		20.930	1.304		46.715	7.725

Values represent the percentage of information dropped when each variable is omitted in each model (the highest values are in bold).

the habitat variables above mentioned remained as the most relevant (even if their contribution to the model diminished in all cases), excluding any climate factors. In winter, after the percentage of dry crops (again the main variable in the model), the amount of wasteland was the most important predictor in the habitat model, while seasonal mean rainfall and net radiation were the most relevant variables in the climate model. The integrated winter models accounted for mixed-selection criteria, based on both habitat and climate variables; while spring models were based mainly in habitat variables.

3.2. Spatial niche models and their width variation

Fig. 2 presents a geographical representation of the spatial niche and its seasonal variation as predicted by MaxEnt. In winter, little bustards disappeared from higher altitude areas, such as foot-hill pastures near the mountains or the high plateau of the eastern and southeastern rim of the region, instead occupying patches along the main river valleys (Jarama and Tagus). Moreover, the winter species distribution in the extensive cereal farmlands of the northeast (most of them included in the Special Protection Area of the Jarama and Henares valleys) was reduced, compared with spring, while it increased in the southern farmlands. According to the habitat models, predicted distribution (associated to niche dimensions) was slightly broader in winter than in spring. However, regarding climate models, conditions were better for the species all across the study area in spring and, therefore, the species was predicted to be more widespread. Boolean integrated seasonal models allowed to compute a slightly larger suitable area (associated to a wider niche) in winter (1058 km²) than in spring (926 km²) (Table 5). This table also presents the range of environmental predictors within suitable patches (removing the effect of the different sampling size on the input data) as a measure of the seasonal change in the niche width. These values again corroborated the general trend in the niche to be larger in winter than in spring. Note that birds supported similar town densities in both seasons, but could be found in areas with a higher proportion of

roads in winter, when they tolerate more human disturbance. However, some land cover percentage values were small due to their reduced availability.

3.3. Changes in landscape connectedness among suitable patches

In the integrated models (based on both habitat and climate) the analysis of connectedness at different thresholds (Fig. 3) showed that: (1) in all cases, suitable patches for the species were more connected across spatial scales in winter; (2) when the threshold was increased (becoming more restrictive), connectedness decreased, more so for spring than for winter; (3) the three indices studied varied in parallel and linear manner for both seasons across scales, with the most similar responses for CLUMPY and PLADJ (both types of contagion indices). These comparable tendencies across thresholds indicate that the results for different indices varied in a proportional way. Therefore, the choice of a particular cut-off had similar effects on different indices.

3.4. Prediction of suitable wintering areas from breeding patches

The inclusion of the spring (breeding) model as a predictor in the integrated full winter model only slightly improved its performance (AUC value increased from 0.940 to 0.948 when this variable was considered), but made spatial patterns more realistic. More relevant was the fact that the spring distribution became the second most important variable (similar to the percentage of wastelands) in this model, behind winter rainfall and surpassing the importance of the amount of dry crops. On the other hand, the inclusion of wintering areas in the spring model also enhanced a bit its performance, although the increase was even smaller (from 0.945 to 0.950). In this case, winter distribution became a less important predictor, being ranked fourth, after dry crops, altitude and pasturelands. These results were consistent with the outcomes achieved when we repeated the models including




Fig. 2 – The figure shows habitat, climate and integrated models for spring and winter (values are cumulated probabilities represented by means of a palette from blue to red). Next, it includes Boolean maps of suitable areas obtained by applying a particular threshold (value of cumulate probability corresponding to a omission error of 10%) on the integrated models (palette black and white). Finally, those Boolean maps are combined to show the overlapping between winter and spring areas.

Table 5 – Percentiles of the environmental variables (i.e. niche dimensions) characterizing the predicted area suitable for the species obtained by applying a threshold of 10% to the full, integrated models (the highest values are in bold)

	Spring ^a				Winter ^b			
	P25	P50	P75	Inter-quartile range (P75 – P25)	P25	P50	P75	Inter-quartile range (P75 – P25)
MDT	626	700	700	74	600	630	700	100
TOPO10	0.00	0.04	0.12	0.12	0.00	0.00	0.08	0.08
TOWNDENS	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
ROADDENS	0.00	0.00	0.16	0.16	0.00	0.00	0.20	0.20
IRRIGPC	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
DRYPERC	0.76	0.92	1.00	0.24	0.68	0.88	1.00	0.32
PASTPERC	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
OLIPERC	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
WASTEPC	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
EVP	27.80	28.70	29.30	1.5	5.65	5.88	6.12	0.47
RAD	50.20	51.00	52.00	1.8	9.96	10.30	10.60	0.64
RAINW	8.33	8.89	9.44	1.11	16.40	17.00	18.20	1.8

^a Available surface: 926 km².
^b Available surface: 1058 km².

507 spring/winter distribution alone as the only variable explain-
 508 ing seasonal distribution: the spring pattern itself was a more
 509 relevant predictor for explaining wintering areas (AUC=0.836)
 510 than vice versa (AUC=0.769).

511 The extensive overlap (591 km²) between spatial niches in
 512 spring and winter, shown in Fig. 3, indicates that potential
 513 areas relevant only for breeding occupied 335 km², while only-
 514 wintering potential areas covered a larger area (467 km²).

4. Discussion

4.1. Can we use MaxEnt to predict seasonal changes in little bustard distribution, environmental preferences and spatial niche width?

518 Although MaxEnt models do not predict the actual limits of
 519 a species' range, they can identify regions with similar envi-
 520 ronmental conditions to occurrence localities (Pearson et al.,
 521 2007). Models presented here provided a good approxima-
 522 tion to little bustard distribution in the region of study, on
 523 the basis of climatic and habitat variables associated with
 524 the species' presence during either spring or the winter.
 525 This result supports MaxEnt as an efficient tool to model
 526 species' occurrence when only small data sets are available,
 527 as already highlighted by previous authors (Elith et al., 2006;
 528 Hernández et al., 2006). On the other hand, ecological niche
 529 theory (Hutchinson, 1957; Chase and Leibold, 2003) predicts
 530 that a species' distribution should be largely determined by its
 531 specific environmental requirements and their spatial varia-
 532 tion (Rosenzweig, 1987). In this sense, our models based
 533 on climatic conditions can be interpreted as the little bustard's
 534 potential distribution range within the region of study, based
 535 on simultaneous variations along different axes of the species'
 536 fundamental niche (Suárez-Seoane et al., 2004). Parallely, the
 537 introduction of variables measuring human disturbances (i.e.
 538 town and road densities) in the habitat models, as a means
 539 of incorporating the species behavioural response (avoidance,
 540 attraction or neutrality) to strong landscape transformations,
 541 bring such models close to be interpreted as realized spatial
 542 niche models (Hutchinson, 1957). However, biotic interactions
 543 (i.e. competition) should be fully measured to obtain more
 544 realistic results. Thus the seasonal shift in the observed niche
 545 could result either from changes in the species' fundamen-
 546 tal niche or from changes in the realized niche, as caused
 547 by the effect of those interactions (Broennimann et al., 2007).
 548 This framework may be used to assess seasonal variations
 549 in the species' habitat preferences and width niche through




Fig. 3 – Variations in the three connectedness indices according to the three different thresholds used to create the Boolean maps.

changes in its distribution ranges. Leaving apart the effect of biotic interactions, birds may respond to environmental stress by relaxing their requirements for habitat selection and thus widening their spatial niche, which allow them to expand their distribution range within the geographic limits imposed by the climatic niche. This was also observed in resident areas, where little bustard's potential suitable habitat slightly increased from winter to spring, even if this difference was lower than expected, probably because the seasonal loss of environmental suitability associated to certain niche factors can be compensated by positive changes in others.

Wintering niche models were, in general terms, less explanatory, predictable and complex than the spring ones, suggesting several facts. First of all, winter distribution was controlled by a smaller number of mixed environmental factors (related to both habitat and climate) than in spring, when birds are breeding and therefore show stricter environmental preferences (mainly related to habitat). This fact is mainly associated to the lower suitability of winter climate conditions in high altitude areas, although some differences in land management, such as the presence in river valleys of more permanent cultures (e.g. irrigated legume crops) or the more frequent set-aside land (e.g. fallows, wastelands) in extensive agricultural areas may also be important (see García de la Morena, 2002; Silva et al., 2004 for the species' winter habitat preferences). In this context, the high winter potential of southern farmlands and a considerable proportion of north-eastern farmlands may reflect the joint effects of climate and land management. In those sectors, little bustards tend to select winter stubble (unpubl. data), a habitat where environmental stress is reduced due to high radiation levels and increased biomass content (weeds and germinated unharvested cereal seeds; Suárez et al., 2004). On the other hand, our results revealed that winter habitat was less predictable than spring, finding the opposite situation for climate conditions. This fact may be related, according to Brotons et al. (2004), to a shift in the restriction of the ecological requirements of the species, which become more generalist in winter, and therefore more difficult to predict.

Spring models were, as before mentioned, controlled by more environmental factors, which defined a narrower niche than in winter, a fact obviously related to strong biotic interactions, reproductive behavior and diet specialization. In this sense, Jiguet et al. (2000) explain that food availability is not critical for little bustard mating selection, as expected for species breeding in exploded leks, although Traba et al. (2008) conclude it is related to display site selection. Those authors also found that large carabids and, in general, large beetles played a relevant role as food resources for males and perhaps for females during the mating season. In this context, considering other niche dimensions (Peterson, 2007) and including demographic parameters into the modeling framework, the reproductive niche would be more realistic and even narrower than suggested by us, as founded by Titeux et al. (2007) for red-backed shrike.

At this point, it must be emphasized that despite the fact that distribution modelling (as developed here) provides relevant insights on the broad-scale environmental niche of species (grinellian models) for conservation and biogeographical research, there are still important conceptual uncertainties

which must be investigated, such as the identification of causal relationships between species distribution and predictors, and the effect of species' ecological characteristics on the performance of the models (McPherson and Jetz, 2007; Tsoar et al., 2007). In this context, besides the development of new algorithms, niche modeling is still hampered by inadequate consideration of critical ecological traits of species, such as dispersal and metapopulation dynamics (Araújo and Guisan, 2006), reproductive parameters (Titeux et al., 2007), biotic interactions and disturbance regime. Also a good knowledge on the ecology of the species is critical to adequately interpret models (Austin, 2002, 2007).

4.2. Can we predict suitable wintering habitat from breeding habitat patches?

The overlap between spring and winter potential distribution areas predicted by our models was high. In fact, most of the identified wintering areas coincided or showed high connectedness with spring patches. Conversely, however, there were significant breeding patches predicted as basically unsuitable which showed little or no connectedness with potential winter areas (foot-hill pastures, eastern rim highlands). We have shown that spring habitat was a much better predictor of the little bustard's wintering range than vice versa, which has clear management implications. For example, preserving breeding sites closer to wintering areas will ensure the conservation of a larger proportion of the total distribution range of the species. Similarly, preserving the largest possible proportion of the potential breeding range will also protect a higher proportion of the wintering range. This is a clear example of how predictive large scale modeling procedures can contribute to the optimization of land management aimed at species conservation. Moreover, we found that fragmentation was perceived differently across the seasons, i.e. suitable patches for the little bustard were more connected across the scales (different thresholds) in winter than in spring, which may have time-dependent implications for species' movement across the landscape. Therefore, we must preserve not only suitable patches, but also agricultural matrix quality (or restore it through, for example, farming extensification) to guarantee dispersion (Donald and Evans, 2006) and enhance its biological connectivity through the year. In particular, for species with seasonally variable food requirements, habitat complementarity may be required to permit movements between these patches (Henle et al., 2004; Donald, 2005) and therefore migratory connectivity across time. This appears to be a useful tool for conservation and management purposes.

5. Conclusions

The applied methodology has allowed us to model successfully little bustard seasonal distribution starting from two input datasets with different sample sizes. Analogously, we have shown how these models can be interpreted as a representation of the species spatial ecological niche at large scale, and may be used to explore niche shifts over a species' annual cycle and their probable causes. In this sense, it must be noticed that even if climate models were only partial (we

may not include all the important variables), they have been highly valuable, particularly when they have been integrated with habitat data. MaxEnt models have proved its efficacy in predicting little bustard winter occurrence from known patches of breeding habitat, which has clear conservation impact in this species, since allows identification of potential winter range and its consideration in large scale management. Additionally, the different relative importance of the areas predicted for each season may help allocating conservation efforts according to the species environmental requirements over the annual cycle.

Acknowledgements

We sincerely wish to thank Andries Rosema, from the EARS (Delft, The Netherlands), for providing EWBSM Meteosat satellite data. Spring censuses were funded by Autonomous Government of Madrid through a contract with SEO-Birdlife. Winter censuses were financed by the project REN 2000-0765 from the Spanish Ministry of Education. We also are grateful to Althea Davies for her help in improving the language and Pilar García who helped in building the DTM in INDUROT (Universidad de Oviedo). María Romero provided her personal vehicle for fieldwork to Manuel B. Morales. Finally, Eladio L. García was funded by the Ministry of Education's FPU Program. The journal editor and an anonymous referee helped with their comments to improve the manuscript.

REFERENCES

- Acevedo, P., Alzaga, V., Cassinello, J., Gortázar, C., 2007. Habitat suitability modelling reveals a strong niche overlap between two poorly known species, the broom hare and the Pyrenean grey partridge in the north of Spain. *Acta Oecol.* **31**, 174–184.
- Aebischer, N.J., Robertson, P.A., Kenward, R.E., 1993. Compositional analysis of habitat use from animal radio-tracking data. *Ecology* **74**, 1313–1325.
- Alonso, J.A., Martín, E., Alonso, J.C., Morales, M.B., Lane, S., 2001. Seasonal movements of male Great Bustards in Spain. *J. Field Ornithol.* **72**, 501–508.
- Araújo, M.B., Guisan, A., 2006. Five (or so) challenges for species distribution modelling. *J. Biogeogr.* **33**, 1677–1688.
- Austin, M.P., 2002. Spatial prediction of species distribution: an interface between ecological theory and statistical modeling. *Ecol. Model.* **157**, 101–118.
- Austin, M.P., 2007. Species distribution models and ecological theory: a critical assessment and some possible new approaches. *Ecol. Model.* **200**, 1–19.
- Beck, J.B., Shultz, E.K., 1986. The use of relative operating characteristic (ROC) curves in test performance evaluation. *Arch. Pathol. Lab. Med.* **110**, 13–20.
- Bibby, C., Burgess, N., Hill, D., Mustoe, S., 2000. *Bird Census Techniques*. Academic Press.
- BirdLife International, 2004. *Birds in Europe II: Population Estimates, Trends and Conservation Status*. BirdLife International, Cambridge, UK.
- Blake, J.G., Loiselle, B.A., 1991. Variation in resource abundance affects capture rates of birds in three lowland habitats in Costa Rica. *The Auk* **108**, 114–130.
- Bolnick, D.I., Svanbäck, R., Araújo, M.S., Persson, L., 2007. Comparative support for the niche variation hypothesis that more generalized populations also are more heterogeneous. *PNAS* **104**, 10075–10079.
- Broennimann, O., Treier, U.A., Müller-Schärer, H., Thuiller, W., Peterson, A.T., Guisan, A., 2007. Evidence of climatic niche shift during biological invasion. *Ecol. Lett.* **10**, 701–709.
- Brooker, L., Brooker, M., Cale, P., 1999. Animal dispersal in fragmented habitat: measuring habitat connectivity, corridor use and dispersal mortality. *Conserv. Ecol.* **3**, 4.
- Brotons, Ll., Thuiller, W., Araújo, M.B., Hirzel, A.H., 2004. Presence-absence versus presence-only modelling methods for predicting bird habitat suitability. *Ecography* **27**, 437–448.
- CAM, 1998. Plan Regional de Estrategia Territorial de la Comunidad de Madrid. Dirección General de Urbanismo y Planificación Territorial, Comunidad de Madrid.
- Chase, J.M., Leibold, M.A., 2003. *Ecological Niches: Linking Classical and Contemporary Approaches*. University of Chicago Press, Chicago.
- Collar, N.J., Crosby, M.J., Stattersfield, A.J., 1994. *Birds to Watch 2: The World List of Threatened Birds*. BirdLife International, Cambridge.
- Cramp, S., Simmons, K.E.L., 1980. *The Birds of the Western Palearctic*. Oxford University Press, London.
- Del Hoyo, J., Elliot, A., Sargatal, J., 1996. *Handbook of the Birds of the World*. Lynx Ed, Barcelona.
- Díaz, M., Martí, R., Gómez-Manzanque, A., Sánchez, A.E., 1994. Atlas de las aves nidificantes en Madrid. SEO-Agencia de Medio Ambiente de la Comunidad de Madrid.
- Donald, P.F., 2005. Climate change and habitat connectivity. Assessing the need for landscape-scale adaptation for birds in the UK. RSPB Research Report 10. RSPB, Sandy, UK.
- Donald, P.F., Evans, A.D., 2006. Habitat connectivity and matrix restoration. *J. Appl. Ecol.* **43**, 209–218.
- Eastman, J.R., 2003. *Idrisi Kilimajaro. User's Guide*. Clark Laboratories, Worcester.
- Elith, J., et al., 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* **29**, 129–151.
- Fielding, A.H., Bell, J.F., 1997. A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ. Conserv.* **24**, 38–49.
- García, J., Suárez-Seoane, S., Miguélez, D., Osborne, P.E., Zumalacárregui, C., 2007. Spatial analysis of the habitat quality in a fragmented population of little bustard. Implications for conservation. *Biol. Conserv.* **137**, 45–56.
- García de la Morena, E.L., 2002. Sisón Común (*Tetrax tetrax*). In: Del Moral, J.C., Molina, B., De la Puente, J., Pérez-Tris, J. (Eds.), Atlas de las aves invernantes de la Comunidad de Madrid. SEO/Birdlife-Monticola, Madrid, pp. 142–143.
- García de la Morena, E.L., Morales, M.B., García, J.T., 2001. Análisis de la importancia de la ZEPA n° 139. Estepas cerealistas de los ríos Jarama y Henares para el sisón común, en el conjunto de su población en la Comunidad de Madrid. Unpublished report. SEO/Birdlife.
- García de la Morena, E.L., Bota, G., Ponjoan, A., Morales, M.B., 2006. El sisón común en España. I censo Nacional. SEO/BirdLife, Madrid.
- García de la Morena, E.L., Morales, M.B., De Juana, E., Suárez, F., 2004a. Does Spanish little bustard migrate? New data on long distance movements of the species. In: Proceedings of the International Symposium on Ecology and Conservation of Steppe-land Birds, Lleida.
- García de la Morena, E.L., Morales, M.B., De Juana, E., Suárez, F., 2007. Surveys of wintering little bustards *Tetrax tetrax* in central Spain: distribution and population estimates at regional scale. *Bird Conserv. Intern.* **17**, 23–34.
- García de la Morena, E.L., De Juana, E., Martínez, C., Morales, M.B., Suárez, F., 2004b. Sisón Común *Tetrax tetrax*. In: Madroño, A., González, C., Atienza, J.C. (Eds.), Libro Rojo de las Aves de España. Dirección General Biodiversidad-SEO/Birdlife, Madrid, pp. 202–207.

- Goriup, P.D., 1994. Little **bustard** *Tetrax tetrax*. In: Tucker, G.M., Heath, M.F. (Eds.), **Birds in Europe: Their Conservation Status**. BirdLife International, Cambridge, pp. 236–237.
- Harris, L.D., 1984. **The Fragmented Forest: Island Biogeography Theory and The Preservation of Biotic Diversity**. University of Chicago Press, Chicago.
- Henle, K., Davis, K.F., Kleyer, M., Margules, C., Settele, J., 2004. Predictors of species sensitivity to fragmentation. **Biodivers. Conserv.** 13, 207–251.
- Hernández, P.A., Graham, C.H., Master, L.L., Albert, D.L., 2006. The effect of sample size and species characteristics on performance of different species distribution modeling methods. **Ecography** 29, 773–785.
- Hutchinson, G.E., 1957. Concluding remarks. Population studies: animal ecology and demography. **Cold Spring Harb. Symp. Quant. Biol.** 22, 415–427.
- Jiguet, F., 2001. Défense des ressources, chix du partenaire et mécanismes de formation des leks chez L'outarde canepetière (*Tetrax tetrax*), une espèce menacée des plaines cerealieres. Université de Paris.
- Jiguet, F., Arroyo, B., Bretagnole, V., 2000. Lek mating systems: a case study in the little bustard *Tetrax tetrax*. **Behav. Proc.** 51, 63–82.
- Leitão, D., Costa, L.T., 2001. First approach to the study of the non-breeding abundance and habitat use by the little bustard *Tetrax tetrax* in the lower Tejo grassland (South Portugal). **Airo** 11, 37–43.
- Martínez, C., 1994. Habitat selection by the Little Bustard *Tetrax tetrax* in cultivated areas of Central Spain. **Biol. Conserv.** 67, 125–128.
- Martínez, C., 1998. Selección de microhábitat del Sisón Común *Tetrax tetrax* durante la estación reproductiva. **Ardeola** 45, 73–76.
- McGarigal, K., Marks, B.J., 1995. FRAGSTATS: Spatial pattern analysis program for quantifying landscape structure. USDA For. Serv. Gen. Tech. Rep. PNW-351.
- McPherson, J.M., Jetz, W., 2007. Effects of species' ecology on the accuracy of distribution models. **Ecography** 30, 135–151.
- Morales, M.B., Alonso, J.C., Alonso, J.A., 2002. Productivity and female individual breeding success in a great bustard *Otis tarda* population. **Ibis** 144, 285–283.
- Morales, M.B., García, J.T., Arroyo, B., 2005. Can landscape composition changes predict spatial and annual variation of little bustard male abundance? **Anim. Conserv.** 8, 167–174.
- Morales, M.B., Alonso, J.C., Alonso, J.A., Martín, E., 2000. Migration patterns in male great bustards (*Otis tarda*). **The Auk** 117, 493–498.
- Q2 Morales, M.B., Traba, J., Carriles, E., Delgado, M.P., García de la **Morena, E.L., in press**. Sexual differences in microhabitat selection of breeding Little Bustards *Tetrax tetrax*: ecological segregation based on vegetation structure. **Acta Oecol.**
- Morin, X., Chuine, I., 2006. Niche breadth, competitive strength and range size of tree species: a trade-off based framework to understand species distribution. **Ecol. Lett.** 9, 185–195.
- Newton, I., 1998. **Population Limitations in Birds**. Academic Press, London.
- Osborne, P.E., Suárez-Seoane, S., 2002. Should data be partitioned spatially before building large-scale distribution models? **Ecol. Model.** 157, 249–259.
- Pain, D.J., Pienkowski, M.W., 1997. **Farming and Birds in Europe**. Academic Press, London.
- Pearce, J.L., Boyce, M.S., 2006. Modelling distribution and abundance with presence-only data. **J. Appl. Ecol.** 43, 405–412.
- Pearson, R.G., Raxworthy, C.J., Nakamura, M., Peterson, A.T., 2007. Predicting species distribution from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. **J. Biogeogr.** 34, 102–117.
- Peterson, A.T., 2007. Why and whywhere: **the** need for more complex models of simpler environmental spaces. **Ecol. Model.** 203, 527–530.
- Phillips, S.J., 2005. Maxent software for species distribution modeling <<http://www.cs.princeton.edu/~schapire/maxent/>>.
- Phillips, S.J., 2008. Transferability, sample selection bias and background data in presence-only modelling: a response to Peterson et al. (2007). **Ecography** 31, 272–278.
- Phillips, S.J., Anderson, R.P., Schapire, R.E., 2006. Maximum entropy modeling of species geographic distributions. **Ecol. Model.** 190, 231–259.
- Rappole, J.H., McDonald, M.V., 1994. Cause and effect in population declines of migratory birds. **Auk** 111, 652–600.
- Robinson, R.A., Sutherland, W.J., 2002. Changes in arable farming and biodiversity in Great Britain. **J. Appl. Ecol.** 39, 157–176.
- Robinson, S.K., Thomson, F.R., Donovan, T.M., Whitehead, D.R., Faaborg, J., 1995. Regional forest fragmentation and the nesting success of migratory birds. **Science** 267, 1987–1990.
- Rosema, A., 1993. Using Meteosat for operational evapotranspiration and biomass monitoring in the Sahel region. **Rem. Sens. Environ.** 45, 1–25.
- Rosema, A., et al., 2001. European energy and water balance monitoring system. EU 4th Framework Program, Contract ENV4-CT97-0478, Final report.
- Rosenzweig, M.L., 1987. Community organization from the point of view of habitat selectors. In: Gee, J.H.R., Giller, P.S. (Eds.), **Organization of Communities: Past and Present**. Blackwell, England, pp. 465–490.
- Rounsevell, M.D.A., Annetts, J.E., Audsley, E., Mayr, T., Reginster, I., 2003. Modelling the spatial distribution of agricultural land use at the regional scale. **Agricult. Ecosyst. Environ.** 95, 465–479.
- Rubenstein, D.R., Hobson, K.A., 2004. From birds to butterflies: animal movement patterns and stable isotopes. **TREE** 19, 256–263.
- Salamolard, M., Moreau, C., 1999. Habitat selection by Little Bustard in a cultivated area of France. **Bird Study** 46, 25–33.
- Sanderson, F.J., Donald, P.F., Burfield, I.J., 2005. Farmland birds in Europe. From policy change to population decline and back again. In: Bota, G., Morales, M.B., Mañosa, S., Camprodon, J. (Eds.), **Ecology and Conservation of Steppe-land Birds**. Lynx Edicions, Barcelona, pp. 211–236.
- Schulz, H., 1985. On the social behaviour of the little bustard: a preliminary report. **Bustard Stud.** 2, 179–181.
- Sherry, T.W., Holmes, R.T., 1996. Winter habitat quality, population limitation and conservation of **Neotropical-Nearctic** migrant birds. **Ecology** 77, 36–48.
- Shirley, S., 2006. Movement of forest birds across river and clearcut edges of varying riparian buffer strip widths. **For. Ecol. Manage.** 223, 190–199.
- Silva, J.P., Pinto, M.A., Palmeirim, J.M., 2004. Managing landscapes for the little bustard *Tetrax tetrax*: lessons from the study of winter habitat selection. **Biol. Conserv.** 117, 521–528.
- Smith, J.N.M., Hellmann, J.J., 2002. Population persistence in fragmented landscapes. **TREE** 17, 397–399.
- Soberon, J., 2007. Grinnellian and eltonian niches and geographic distributions of species. **Ecol. Lett.** 10, 1115–1123.
- Suárez, F., Naveso, M.A., De Juana, E., 1997. Farming in drylands of Spain: birds of the pseudosteppes. In: Pain, D.J., Pienkowski, M.W. (Eds.), **Farming and Birds in Europe**. Academic Press, London, pp. 297–330.
- Suárez, F., Garza, V., Oñate, J.J., García de la Morena, E.L., Ramírez, A., Morales, M.B., 2004. Adequacy of winter stubble maintenance for steppe passerine conservation in central Spain. **Agricult. Ecosyst. Environ.** 104, 667–671.
- Suárez-Seoane, S., Osborne, P.E., Alonso, J.C., 2002. Large-scale habitat selection by agricultural steppe birds in Spain: identifying species-habitat responses using Generalised Additive Models. **J. Appl. Ecol.** 39, 755–777.

- 929 Suárez-Seoane, S., Osborne, P.E., Rosema, A., 2004. Can climate
930 data from METEOSAT improve wildlife distribution models?
931 *Ecography* 27, 629–636.
- 932 Tellería, J.L., Suárez, F., Santos, T., 1988. Bird communities of the
933 Iberian shrubsteppes. *Holarctic Ecol.* 11, 171–177.
- 934 Thuiller, W., Richardson, D.M., Pysek, P., Midgley, G.F., Hughes,
935 G.O., Rouget, M., 2005. Niche-based modeling as a tool for
936 predicting the risk of alien plant invasions at a global scale.
937 *Global Change Biol.* 11, 2234–2250.
- 938 Titeux, N., Dufrené, M., Radoux, J., Hirzel, A.H., Defourny, P., 2007.
939 Fitness-related parameters improve presence-only
940 distribution modelling for conservation practice: the
941 case of the red-backed shrike. *Biol. Conserv.* 138,
942 207–223.
- 943 Traba, J., Morales, M.B., García de la Morena, E.L., Delgado, M.P.,
944 2008. Selection of breeding territory by little bustard (*Tetrax*
945 *tetrax*) males in Central Spain: the role of arthropod
946 availability. *Ecol. Res.*, doi:10.1007/s11284-007-0418-4.
- 947 Tsoar, A., Allouche, O., Steinitz, O., Rotem, D., Kadman, R., 2007. A
948 comparative evaluation of presence-only methods for
949 modeling species distribution. *Div. Distrib.* 13,
397–405.
- Turchin, P., 1991. Translating foraging movements in
950 heterogeneous environments into the spatial distribution of
951 foragers. *Ecology* 72, 1253–1266.
- 952 Van Valen, L., 1965. Morphological variation and width of
953 ecological niche. *Am. Nat.* 99, 377–389.
- 954 Wiens, J.A., 1989. *The Ecology of Bird Communities*. Cambridge
955 University Press.
- 956 Wolff, A., 2001. Changements agricoles et conservation de la
957 grande avifaune de plaine: Etude des relations
958 espèce-habitats à différentes échelles chez l'Outarde
959 canepetière. Université Montpellier II, Université des Sciences
960 et Techniques du Languedoc.
- 961 Wolff, A., Dieuleveut, T., Martin, J.L., Bretagnolle, V., 2002.
962 Landscape context and little bustard abundance in a
963 fragmented steppe: implications for reserve management in
964 mosaic landscapes. *Biol. Conserv.* 107, 211–220.
- 965 Wolff, A., Paul, J.P., Martin, J.L., Bretagnolle, V., 2001. The benefits
966 of extensive agriculture to birds: the case of the little bustard.
967 *J. Appl. Ecol.* 38, 963–975.
- 968 Zweig, M.H., Campbell, G., 1993. Receiver-operating
969 characteristics (ROC) plots: a fundamental evaluation tool in
970 clinical medicine. *Clin. Chem.* 39, 561–577.
- 971