

SUPPLEMENTARY DATA

Table 1 of the supplementary data. Standardized mean differences before (raw) and after (matched) matching

	Standardized mean difference (raw)	Standardized mean difference (matched)
Age	0.7214	0.0105
Age > 80 years	0.4256	0.0118
Diabetes mellitus ^a	0.1414	-0.0063
Hypertension ^b	0.1797	0.0859
Dyslipidemia ^c	0.0049	0.0490
Obesity ^d	0.2413	0.0187
Chronic kidney failure ^e	0.0686	-0.0218
Smoking habit (current or former)	-1.0393	-0.0119
Cardiogenic shock at presentation	0.1279	-0.0078
Oral intubation	0.0615	0.0620
Cardiac resuscitation at presentation	0.1279	-0.0078
Left ventricular ejection fraction < 35%	0.1554	0.0301
anterior wall infarct location	0.0062	-0.0862
Femoral access	0.0900	0.0076
Use of GP IIb-IIIa	-0.2464	-0.0268
Bivalirudin	0.0797	0.0555
Aortic counterpulsation	0.0617	-0.0452
Thrombus aspiration	-0.1952	-0.0105
Stent implantation		
Drug eluting stent implantation	-0.0881	0.0506

^a American Heart Association Guidelines definition

^b Arterial systemic pressure \geq 140/90 mmHg.

^c Total cholesterol \geq 200 mg/dL; low density lipoprotein \geq 130 mg/dL; triacylglycerol \geq 175 mg/dL.

^d Body mass index > 30

^e Dialysis or serum creatinine > 2 mg/dL

Table 2. Main outcome analysis stratifying matched population according to 2 time-period

	2006 - 2010			2011 - 2016		
	Female n = 175	Male n = 175	<i>P</i>	Female (n = 204)	Male (n = 204)	<i>P</i>
Procedural efficacy	165 (94.3)	168 (94.3)	.999	197 (96.6)	191 (95.0)	.468
Long term BARC bleeding	8 (1.7)	3 (4.6)	.137	5 (2.5)	4 (2.0)	.999
Total mortality at long-term follow-up	62 (35.4)	75 (42.1)	.230	47 (23.0)	50 (24.9)	.727

BARC, Bleeding Academic Research Consortium.