




Melatonin affects red deer spermatozoa motility and physiology in capacitating and non-capacitating conditions

Estela Fernández-Alegre¹ | Estíbaliz Lacalle¹ | Cristina Soriano-Úbeda^{2,3}  |
Juan Carlos Domínguez^{2,4} | Adriana Casao⁵  | Felipe Martínez-Pastor^{2,3} 

¹Bianor Biotech SL, León, Spain

²Institute for Animal Health and Cattle Development (INDEGSAL) and IMAPOR Research Group, Universidad de León, León, Spain

³Department of Molecular Biology (Cell Biology), Universidad de León, León, Spain

⁴Department of Animal Medicine, Surgery and Anatomy (Animal Medicine and Surgery), University of León, León, Spain

⁵Department of Biochemistry and Molecular and Cell Biology, Institute of Environmental Sciences of Aragón, School of Veterinary Medicine, University of Zaragoza, Zaragoza, Spain

Correspondence

Cristina Soriano-Úbeda, INDEGSAL and IMAPOR, Universidad de León, León, Spain.

Email: c.soriano.ubeda@unileon.es

Funding information

MINECO, Grant/Award Number: AGL2013-43328

Abstract

Melatonin affects sperm physiology, possibly through membrane receptors. Effects were tested at low concentrations (1 pM, 100 pM, 10 nM and 1 µM) in red deer epididymal spermatozoa as a model for high-seasonality species. Samples were incubated with melatonin as uncapacitated or capacitating conditions (heparin) and evaluated for motility and physiology (flow cytometry). Most effects occurred at low concentrations (nM-pM), mainly protecting from apoptosis and maintaining acrosomal integrity, suggesting a role for membrane receptors rather than a direct antioxidant effect. Intracellular calcium was not affected, differing from other studies and perhaps because of the epididymal origin. This study supports the relevance of melatonin on sperm physiology and could contribute to the application of reproductive technologies in wild ruminants.

KEYWORDS

melatonin, red deer, sperm capacitation, sperm physiology, spermatozoon

1 | INTRODUCTION

Melatonin is a widespread and pleiotropic molecule (Hardeland et al., 2011). Widely studied as a hormone regulating the circadian cycle and seasonality, evidence has mounted on its paracrine activity in reproductive organs (Gonzalez-Arto et al., 2016; Martínez-Marcos et al., 2019) and its membrane receptors MT₁ and MT₂ are present in spermatozoa (González-Arto et al., 2016, 2017). Melatonin affects sperm survival, capacitation and chemotaxis (Casao et al., 2012; Cebrián-Pérez et al., 2014; Espino et al., 2011). This topic has been studied in some domestic ruminants (Fernández-Alegre et al., 2020; González-Arto et al., 2016), but it could be different for wild ruminants, subjected to pronounced seasonal reproductive rhythms (Martínez-Pastor et al., 2005). Our objective was to test its role on

red deer (*Cervus elaphus*) spermatozoa under non-capacitating or capacitating conditions, with the hypothesis that it could affect them at concentrations compatible with receptor activation, as reported for bull (Fernández-Alegre et al., 2020).

2 | MATERIALS AND METHODS

The experimental design and methods followed our study on bull (Fernández-Alegre et al., 2020) and are expanded in the supplementary material. Spermatozoa were obtained by cuts from the cauda epididymis of six adult deers (testicles harvested after regulated hunting at Riaño reserve and Doñana National Park, Spain) (Martínez-Pastor et al., 2006) and extended in TALP-HEPES at

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Reproduction in Domestic Animals* published by Wiley-VCH GmbH

TABLE 1 Effect of different melatonin concentrations (1 μ M, 10 nM, 100 pM and 1 pM; CTL as the vehicle-only control, 0.2% DMSO) in non-capacitating and capacitating conditions

Parameter	Non-capacitated					Capacitated				
	CTL	1 μ M	10 nM	100 pM	1 pM	CTL	1 μ M	10 nM	100 pM	1 pM
Total motility (%)	68.7 \pm 3.8	70.1 \pm 3.5	70.9 \pm 2.9	69.0 \pm 1.6	72.2 \pm 4.1	60.0 \pm 2.7	63.3 \pm 2.4	70.0 \pm 2.3**	61.7 \pm 3.7	63.7 \pm 2.7
Progressive motility (%)	34.5 \pm 3.6	39.5 \pm 2.7	38.2 \pm 2.5	34.0 \pm 1.9	35.9 \pm 3.3	30.2 \pm 2.8	30.1 \pm 2	39.4 \pm 2.8**	30.6 \pm 2.7	30.6 \pm 2.2
VAP (μ m/s)	91.7 \pm 5.0	98.5 \pm 2.6	106.2 \pm 1.8**	102.3 \pm 2.6	100.9 \pm 3.8	112.8 \pm 2.2	102.3 \pm 4.0*	109.6 \pm 3.1	104.4 \pm 4.0	104.0 \pm 3.9
STR (%)	80.1 \pm 1.8	82.1 \pm 2.2	80.4 \pm 1.8	74.8 \pm 2.7*	79.1 \pm 1.4	73.0 \pm 2.6	74.0 \pm 2.3	82.9 \pm 3.4*	75.9 \pm 3.0	73.8 \pm 1.9
ALH (μ m)	1.9 \pm 0.1	1.9 \pm 0.1	2.0 \pm 0.0	2.0 \pm 0.0*	2.0 \pm 0.1**	2.3 \pm 0.1	2.1 \pm 0.0	2.1 \pm 0.0	2.2 \pm 0.1	2.2 \pm 0.1
Viability (%; PI ⁻)	78.7 \pm 0.6	77.9 \pm 0.7	77.1 \pm 1.1	78.6 \pm 0.9	80.1 \pm 0.5	79.2 \pm 0.8	77.4 \pm 0.8	75 \pm 1.4**	75.7 \pm 0.7*	76.1 \pm 1.3
Viability (%; YO-PRO-1 ⁻)	58.3 \pm 1.6	60.1 \pm 1.9	57.3 \pm 2.7	62.4 \pm 2.6	58.8 \pm 1.2	45.3 \pm 1.0	51.1 \pm 1.5*	51.8 \pm 2.3*	50.3 \pm 2.9*	54.1 \pm 1.6**
Apoptotic (%; ratio PI ⁻)	23.1 \pm 1.7	19.3 \pm 3.1	21.5 \pm 1.4	14.3 \pm 3.8***	22.6 \pm 1.1	34.4 \pm 2.2	29.5 \pm 3.8	24.7 \pm 2.4**	30.0 \pm 3.2	24.8 \pm 1.9**
Reacted acrosomes (%)	29.6 \pm 0.7	29.7 \pm 0.8	31.2 \pm 0.6	25.4 \pm 1.9*	25.3 \pm 2.2**	34.2 \pm 2.4	35.1 \pm 2.1	34.0 \pm 1.2	36.2 \pm 1.8	31.0 \pm 1.9
Reacted acrosomes (%; ratio PI ⁻)	20.0 \pm 0.7	19.7 \pm 0.5	20.7 \pm 0.3	14.6 \pm 2.6**	14.9 \pm 2.7**	26.5 \pm 2.4	26.3 \pm 2.3	23.7 \pm 1.7	27.4 \pm 2.3	23.0 \pm 2.3
Capacitated (%; ratio PI ⁻)	17.5 \pm 2.4	16.2 \pm 1.4	16.5 \pm 1.9	15.7 \pm 1.3	11.1 \pm 0.8**	17.1 \pm 1.2	15.8 \pm 1.1	16.2 \pm 1.2	17.6 \pm 0.7	15.3 \pm 0.9
Active mitochondria (%)	63.7 \pm 1.8	64.4 \pm 0.9	63.2 \pm 2.5	67.7 \pm 0.8	69.0 \pm 1.5*	55.8 \pm 0.9	59.6 \pm 1.6*	57.4 \pm 1.8	61.9 \pm 0.6**	58.8 \pm 0.7
[Ca ²⁺] _i (MFI)	6.7 \pm 0.1	7.1 \pm 0.2	6.8 \pm 0.2	6.7 \pm 0.2	6.7 \pm 0.2	6.5 \pm 0.1	6.4 \pm 0.1	6.7 \pm 0.2	6.8 \pm 0.1	6.6 \pm 0.1
Cytoplasmic ROS (MFI)	3.9 \pm 0.1	4.0 \pm 0.2	4.0 \pm 0.1	3.9 \pm 0.1	3.7 \pm 0.1	3.7 \pm 0.1	3.8 \pm 0.1	3.8 \pm 0.1	3.9 \pm 0.1	3.8 \pm 0.1
Mitochondrial ROS (%; ratio PI ⁻)	21.1 \pm 1.9	17.1 \pm 0.6	19.1 \pm 2.2	11.5 \pm 2.9**	18.8 \pm 1.7	24.1 \pm 1.3	22.7 \pm 1.8	19.4 \pm 0.6	25.2 \pm 2.9	17.5 \pm 1.9*

Note: Results are shown as mean \pm SEM. Significant effects on the control in each group are indicated by * p < .05, ** p < .01, *** p < .001. Parameters are described in detail in the main text and the supplementary material.

50 × 10⁶/ml. Tubes were split, one receiving 2 U/ml heparin (capacitated) and split again, adding melatonin at 1 μM, 10 nM, 100 pM, 1 pM and control (vehicle, 0.2% DMSO).

The samples were incubated for 4 h (38°C, 5% CO₂) and analysed by CASA for motility (ISAS; Proiser, Valencia, Spain) and flow cytometry. For flow cytometry (CyAn ADP, Beckman Coulter), fluorescent probes were: Hoechst 33,342 (H342), 5 μM; YO-PRO-1 (YP1), 100 nM; Fluo-4 (Fluo4), 100 nM; CM-H₂DCFDA (CFDA) at 5 μM; merocyanine 540 (M540), 2 μM; propidium iodide (PI), 1 μM; PNA-AlexaFluor 647 (PNA), 1 μg/ml; MitoTracker deep red (MT), 100 nM; MitoSOX (MSX), 1 μM. Aliquots were added at 10⁶ ml⁻¹ to combinations of H342/Fluo4/M540/PI/PNA, H342/CFDA/PI and H342/YP/MSX/MT and analysed after 15 min in the dark.

The statistical analysis was performed with R using linear mixed-effects models (treatment and incubation as fixed effects; male as random effect). Results are shown as mean ± SEM.

3 | RESULTS

Table 1 summarizes the results for melatonin effects on the non-capacitated and capacitated groups (further details on the Tables S1 and S2). The proportions of motile spermatozoa (total and progressive) increased in 10 nM only in capacitated samples ($p < 0.01$). Some kinematic variables were significantly affected in non-capacitated samples by nM-pM, suggesting a stimulation, with few effects in the capacitated group.

Sperm viability moderately decreased in capacitated (10 nM and 100 pM, $p < 0.05$), but the proportion of non-apoptotic spermatozoa significantly increased in all cases, with a drop for the apoptotic ratio in 10 nM and 1 pM (and 100 pM in uncapacitated). Acrosomal damage significantly decreased in uncapacitated by pM. Mitochondrial activity increased by 1 pM in uncapacitated and 1 μM and 100 pM in capacitated. [Ca²⁺]_i and cytoplasmic ROS were not affected, but mitochondrial ROS were abated by 100 pM in non-capacitated ($p < 0.01$) and 1 pM in capacitated samples ($p < 0.05$).

4 | DISCUSSION

Melatonin affected red deer epididymal spermatozoa at the low range of concentrations, consistent with our previous study on bull semen (Fernández-Alegre et al., 2020) and exerting a role on sperm physiology through its membrane receptors. In these conditions, a direct antioxidant effect seems unlikely. Our results support previous evidence on a functional role for MT₁ and MT₂ in spermatozoa (Casao et al., 2012; Fujinoki, 2008; González-Arto et al., 2016), consistently showing their implication in sperm capacitation modulation keeping motility and preserving viability and oxidation state.

This first study in red deer spermatozoa hints at a protective effect, possibly by inhibiting apoptosis-related pathways (Casao et al., 2010; Espino et al., 2011). However, we could not detect intracellular calcium changes, contrarily to studies in ejaculated

spermatozoa (Fernández-Alegre et al., 2020; Gimeno-Martos et al., 2019). Epididymal spermatozoa are quiescent, not having contacted seminal plasma (Martínez et al., 2008), and signalling pathways modifying calcium patterns could be inhibited at this stage (Ernesto et al., 2015; Gibbs et al., 2011).

This study suggests differences between species and possibly epididymal and ejaculated spermatozoa and could be helpful for the conservation and application of wild ruminant post-mortem samples.

ACKNOWLEDGEMENTS

The authors thank Dr. Joaquín Vicente and SaBio (IREC, CSIC-UCL-JCMM, Spain) and hunting wardens at Riaño Reserve for providing the samples, and Indira Álvarez-Fernández and Cristina Arija. Supported by grant AGL2013-43328 (MINECO, Spain).

CONFLICT OF INTEREST

None declared.

ORCID

Cristina Soriano-Úbeda  <https://orcid.org/0000-0003-2880-3342>

Adriana Casao  <https://orcid.org/0000-0003-1997-4262>

Felipe Martínez-Pastor  <https://orcid.org/0000-0003-2987-4302>

REFERENCES

- Casao, A., Luna, C., Pérez-Pe, R. N., Muio-Blanco, T., & Cebrián-Pérez, J. A. (2012). Receptor mediated effect of melatonin on ram sperm calcium patterns. *Reproduction in Domestic Animals*, 47, 90. <https://doi.org/10.1111/j.1439-0531.2012.02044.x/full>
- Casao, A., Mendoza, N., Pérez-Pé, R., Grasa, P., Abecia, J.-A., Forcada, F., Cebrián-Pérez, J. A., & Muiño-Blanco, T. (2010). Melatonin prevents capacitation and apoptotic-like changes of ram spermatozoa and increases fertility rate. *Journal of Pineal Research*, 48(1), 39–46. <https://doi.org/10.1111/j.1600-079X.2009.00722.x>
- Cebrián-Pérez, J., Casao, A., González-Arto, M., Dos Santos Hamilton, T., Pérez-Pé, R., & Muiño-Blanco, T. (2014). Melatonin in sperm biology: Breaking paradigms. *Reproduction in Domestic Animals*, 49, 11–21. <https://doi.org/10.1111/rda.12378>
- Ernesto, J. I., Weigel Muñoz, M., Battistone, M. A., Vasen, G., Martínez-López, P., Orta, G., Figueiras-Fierro, D., De la Vega-Beltran, J. L., Moreno, I. A., Guidobaldi, H. A., Giojalas, L., Darszon, A., Cohen, D. J., & Cuasnicú, P. S. (2015). CRISP1 as a novel CatSper regulator that modulates sperm motility and orientation during fertilization. *The Journal of Cell Biology*, 210(7), 1213–1224. <https://doi.org/10.1083/jcb.201412041>
- Espino, J., Ortiz, Á., Bejarano, I., Lozano, G. M., Monllor, F., García, J. F., Rodríguez, A. B., & Pariente, J. A. (2011). Melatonin protects human spermatozoa from apoptosis via melatonin receptor- and extracellular signal-regulated kinase-mediated pathways. *Fertility and Sterility*, 95(7), 2290–2296. <https://doi.org/10.1016/j.fertnstert.2011.03.063>
- Fernández-Alegre, E., Álvarez-Fernández, I., Domínguez, J. C., Casao, A., & Martínez-Pastor, F. (2020). Melatonin non-linearly modulates bull spermatozoa motility and physiology in capacitating and non-capacitating conditions. *International Journal of Molecular Sciences*, 21(8), 2701. <https://doi.org/10.3390/ijms21082701>
- Fujinoki, M. (2008). Melatonin-enhanced hyperactivation of hamster sperm. *Reproduction*, 136(5), 533–541. <https://doi.org/10.1530/REP-08-0202>

- Gibbs, G. M., Orta, G., Reddy, T., Koppers, A. J., Martínez-López, P., Luis de la Vega-Beltrán, J., Lo, J. C. Y., Veldhuis, N., Jamsai, D., McIntyre, P., Darszon, A., & O'Bryan, M. K. (2011). Cysteine-rich secretory protein 4 is an inhibitor of transient receptor potential M8 with a role in establishing sperm function. *Proceedings of the National Academy of Sciences of the United States of America*, 108(17), 7034–7039. <https://doi.org/10.1073/pnas.1015935108>
- Gimeno-Martos, S., Casao, A., Yeste, M., Cebrián-Pérez, J. A., Muiño-Blanco, T., & Pérez-Pé, R. (2019). Melatonin reduces cAMP-stimulated capacitation of ram spermatozoa. *Reproduction, Fertility, and Development*, 31(2), 420–431. <https://doi.org/10.1071/RD18087>
- González-Arto, M., Aguilar, D., Gaspar-Torrubia, E., Gallego, M., Carvajal-Serna, M., Herrera-Marcos, L. V., Serrano-Blesa, E., Hamilton, T. R. D. S., Pérez-Pé, R., Muiño-Blanco, T., Cebrián-Pérez, J. A., & Casao, A. (2017). Melatonin MT₁ and MT₂ receptors in the ram reproductive tract. *International Journal of Molecular Sciences*, 18(3), E662. <https://doi.org/10.3390/ijms18030662>
- González-Arto, M., Hamilton, T. R. D. S., Gallego, M., Gaspar-Torrubia, E., Aguilar, D., Serrano-Blesa, E., Abecia, J. A., Pérez-Pé, R., Muiño-Blanco, T., Cebrián-Pérez, J. A., & Casao, A. (2016). Evidence of melatonin synthesis in the ram reproductive tract. *Andrology*, 4(1), 163–171. <https://doi.org/10.1111/andr.12117>
- González-Arto, M., Vicente-Carrillo, A., Martínez-Pastor, F., Fernández-Alegre, E., Roca, J., Miró, J., Rigau, T., Rodríguez-Gil, J. E., Pérez-Pé, R., Muiño-Blanco, T., Cebrián-Pérez, J. A., & Casao, A. (2016). Melatonin receptors MT₁ and MT₂ are expressed in spermatozoa from several seasonal and nonseasonal breeder species. *Theriogenology*, 86(8), 1958–1968. <https://doi.org/10.1016/j.theriogenology.2016.06.016>
- Hardeland, R., Cardinali, D. P., Srinivasan, V., Spence, D. W., Brown, G. M., & Pandi-Perumal, S. R. (2011). Melatonin—A pleiotropic, orchestrating regulator molecule. *Progress in Neurobiology*, 93(3), 350–384. <https://doi.org/10.1016/j.pneurobio.2010.12.004>
- Martínez, A. F., Martínez-Pastor, F., Alvarez, M., Fernández-Santos, M. R., Esteso, M. C., de Paz, P., Garde, J. J., & Anel, L. (2008). Sperm parameters on Iberian red deer: Electroejaculation and post-mortem collection. *Theriogenology*, 70(2), 216–226. <https://doi.org/10.1016/j.theriogenology.2008.04.001>
- Martínez-Marcos, P., Carvajal-Serna, M., Lázaro-Gaspar, S., Pérez-Pé, R., Muiño-Blanco, T., Cebrián-Pérez, J. A., & Casao, A. (2019). Presence of melatonin-catabolizing non-specific enzymes myeloperoxidase and indoleamine 2,3-dioxygenase in the ram reproductive tract. *Reproduction in Domestic Animals*, 54(12), 1643–1650. <https://doi.org/10.1111/rda.13574>
- Martínez-Pastor, F., Díaz-Corujó, A. R., Anel, E., Herráez, P., Anel, L., & de Paz, P. (2005). Post mortem time and season alter subpopulation characteristics of Iberian red deer epididymal sperm. *Theriogenology*, 64(4), 958–974. <https://doi.org/10.1016/j.theriogenology.2005.01.003>
- Martínez-Pastor, F., García-Macías, V., Álvarez, M., Chamorro, C., Herráez, P., de Paz, P., & Anel, L. (2006). Comparison of two methods for obtaining spermatozoa from the cauda epididymis of Iberian red deer. *Theriogenology*, 65(3), 471–485. <https://doi.org/10.1016/j.theriogenology.2005.05.045>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Fernández-Alegre, E., Lacalle, E., Soriano-Úbeda, C., Carlos Domínguez, J., Casao, A., & Martínez-Pastor, F. (2022). Melatonin affects red deer spermatozoa motility and physiology in capacitating and non-capacitating conditions. *Reproduction in Domestic Animals*, 57(Suppl. 5), 82–85. <https://doi.org/10.1111/rda.14137>